

Publications of Dr. Hiroaki SATO, 2005-2015

- Kanda N, Yokota T, Shibata E, and Sato H (2005) Diversity of dung-beetle community in declining Japanese subalpine forest caused by an increasing sika deer population. *Ecological Research* 20: 135-141.
- Komori R, Sato H, Harumoto T, and Takagi Y (2005) A new mutation in the timing of autogamy in *Paramecium tetraurelia*. *Mechanisms of Ageing and Development* 126: 752-759.
- Katano I, Mitsuhashi H, Isobe Y, Sato H, Ohishi T. (2005) Reach-scale distribution dynamics of a grazing stream insect, *Micrasema quadriloba* Martynov (Trichoptera), in relation to current velocity and periphyton abundance. *Zoological Science* 22: 853-860.
- Ishida TA, Hattori K, Shibata S, Suzuki M, Sato H, Kimura MT (2005) Sex allocation of a cosexual wind-pollinated tree *Quercus dentata* in terms of four currencies. *Journal of Plant Research* 118: 193-197.
- Oishi M, Yokota T, Teramoto N, and Sato H (2006) Feeding preference and performance of the Japanese oak silkmoth for upper-crown and lower-crown leaves. *Entomological Science* 9: 161-169.
- Katano I, Mitsuhashi H, Isobe Y, Sato H, Oishi T (2007) Group size of feeding stream case-bearing caddisfly grazers and resource abundance. *Basic and Applied Ecology* 8: 269-279.
- Kitamura M, Nakamura T, Hattori K, Ishida TA, Shibata S, Sato H, and Kimura MT (2007) Among-tree variation in leaf traits and herbivore attacks in a deciduous oak, *Quercus dentata*. *Scandinavian Journal of Forest Research* 22: 211-218.
- Oishi M and Sato H (2007) Inhibition of premature leaf abscission by a leafminer and its adaptive significance. *Environmental Entomology*, 36: 1504-1511.
- Kato T, Ishida K and Sato H (2008) The evolution of nettle resistance to heavy deer browsing. *Ecological Research*, 23: 339-345.
- Goda R, Ando M, Sato H and Shibata E (2008) Application of fecal pellet group count to sika deer (*Cervus nippon*) population monitoring on Mt. Ohdaigahara, central Japan. *Mammal Study*, 33, 93-97.
- Oishi M and Sato H (2008) Guild Structure and coexistence mechanisms in the parasitoid assemblage associated with a leafminer, *Coptotriche japoniella* (Lepidoptera, Tischeriidae), on an evergreen tree, *Eurya japonica* (Theaceae). *Environmental Entomology*, 37, 1231-1240.
- Nakamura T, Hattori K, Ishida TA, Sato H, and Kimura MT (2008). Population dynamics of

- leafminers on a deciduous oak *Quercus dentata*. *Acta Oecologia*, 34, 259–265.
- Oishi M and Sato H (2009) Life history traits, larval habits and larval morphology of a leafminer, *Coptotriche japoniella* (Tischeriidae), on an evergreen tree, *Eurya japonica* (Theaceae). *Journal of the Lepidopterist's Society*, 63, 93-99.
- Hata Y, Hashiba T, Nakamura T, Kitamura M, Ishida TA, Akimoto S, Sato H and Kimura MT (2011) Differences in leafminer (*Phyllonorycter*, Gracillariidae, Lepidoptera) and aphid (*Tverculatus*, Aphididae, Hemiptera) composition among *Quercus dentata*, *Q. crspula*, *Q. serrata*, and their hybrids. *Journal of Forest Research*, 16, 309-318.
- Sato H (2011) The family Nepticulidae. Komai F, Yoshiyasu H, Nasu Y, Saito H (eds) A guide to the Lepidoptera of Japan, pp. 99-105, 552-553, pl. 3. Tokai University Press, Tokyo (In Japanese)
- Sato H (2011) The superfamily Tischerioidea. Komai F, Yoshiyasu H, Nasu Y, Saito H (eds) A guide to the Lepidoptera of Japan, pp. 126-132, 558-559, pl. 7. Tokai University Press, Tokyo (In Japanese)
- Sato H and Kumata T (2011) The family Gracillariidae. Komai F, Yoshiyasu H, Nasu Y, Saito H (eds) A guide to the Lepidoptera of Japan, pp. 149-159, 568-573, pls.13-16. Tokai University Press, Tokyo (In Japanese)
- Shinozaki Y, Yamamoto Y, Oishi M, and Sato H (2012) Early leaf abscission has little effect on larval mortality of *Ectoedemia cerviparadisicola* (Lepidoptera, Nepticulidae) associated with *Quercus gilva*. *Annals of Entomological Society of America*, 105, 572-578.
- Shikata M, Kato T, Shibata E and Sato H. (2013) Among-population variation in resistance traits of a nettle and its relationship with deer habitat use frequency. *Ecological Research*, 28, 207-216.
- Iwamoto M, Horikawa C, Shikata M, Wasaka N, Kato T, Sato H (2014) Stinging hairs on the Japanese nettle *Urtica thunbergiana* have a defensive function against mammalian but not insect herbivores. *Ecological Research*, 29, 455-462.